

POSTER PRESENTATION

Open Access

Changes in atherosclerotic plaque composition assessed by MRI in the superficial femoral artery with two years of lipid lowering therapy

Amy M West*, AJ Pesch, Neesurg Mehta, Justin D Anderson, Frederick H Epstein, Craig H Meyer, Klaus D Hagspiel, Stuart S Berr, Nancy L Harthun, Jennifer R Hunter, Joseph M DiMaria, John M Christopher, Christopher M Kramer

From 2011 SCMR/Euro CMR Joint Scientific Sessions
Nice, France. 3-6 February 2011

Objective

We hypothesized that changes in atherosclerotic plaque composition with lipid lowering therapy could be measured in patients with peripheral arterial disease (PAD) using MRI of the superficial femoral artery (SFA).

Background

MRI of atherosclerotic plaque components in the carotid arteries correlates with histology, is reproducible, and can monitor lipid lowering therapy.

Methods

43 patients with PAD (age 63 ± 9 , ABI 0.72 ± 0.14) were imaged at baseline and annually for 2 years after beginning lipid lowering therapy. Twenty statin-naïve patients were randomized to simvastatin (S, $n=11$) or simvastatin +ezetimibe (S+E, $n=9$). Patients already on statins not at LDL cholesterol goal had open-label ezetimibe added (E, $n=23$). MRI of a single slice of the SFA was performed on Siemens Avanto 1.5T scanner with a proton-density weighted image (TR 1100ms, TE 7.6ms), T1-W image (TR 700ms, TE 7.7ms) and T2-W image (TR 2000ms, TE 85ms), using slice thickness 3mm, matrix 256x256 pixels, FOV 150mm, flip angle 180°. Image location was repeated over time using the distance of the imaged slice from the bifurcation of the common femoral artery. Image quality scale from 1-4 was used to grade the images, excluding 6 studies for score ≤ 2 in more than one image. VesselMASS software was used to measure plaque components (lipids, calcium, loose matrix and fibrous tissue) based on published criteria for appearance on T1/T2/PDW images. Changes in parameters were

assessed with ANOVA repeated measures model and correlation with two-tailed Pearson.

Results

LDL at baseline in the statin-naïve patients (S and S+E) was 127 ± 34 mg/dl and fell at one year (82 ± 37 , $p < 0.001$) and was unchanged at two years (84 ± 44). There was no difference in LDL between S and S+E. The LDL in E at baseline was 93 ± 21 mg/dl, decreased significantly at one year (77 ± 35 , $p < 0.001$), and was similar at year 2 (74 ± 27). Total cholesterol changes paralleled that of LDL. See Table 1 for changes in wall area and plaque components. The amount of plaque lipids correlated with wall area ($r=0.44$, $p < 0.003$) for all studies. Intra-observer

Table 1 Changes in plaque over time

	Baseline	Year 1	Year 2
Wall Area (cm²)			
S and S+E	0.33 ± 0.14	0.32 ± 0.14	0.32 ± 0.14
E	0.31 ± 0.05	0.29 ± 0.12	0.29 ± 0.12
Lipids (cm²)			
S and S+E	0.05 ± 0.04	0.04 ± 0.04	$0.03 \pm 0.03^*$
E	0.04 ± 0.05	0.04 ± 0.05	$0.02 \pm 0.03^*$
Calcium (cm²)			
S and S+E	0.004 ± 0.007	0.005 ± 0.010	0.009 ± 0.016
E	0.011 ± 0.026	0.011 ± 0.022	0.014 ± 0.023
Loose Matrix (cm²)			
S and S+E	0.05 ± 0.07	0.04 ± 0.05	0.03 ± 0.03
E	0.03 ± 0.06	0.04 ± 0.07	0.04 ± 0.07
Fibrous Tissue (cm²)			
S and S+E	0.23 ± 0.13	0.23 ± 0.12	0.25 ± 0.13
E	0.23 ± 0.09	0.20 ± 0.10	0.22 ± 0.10

*ANOVA $p < 0.05$.

University of Virginia, Charlottesville, VA, USA




Figure 1 Representative PDW (left), T1-W (center) and T2-W (right) images of the SFA (arrow) in a study patient.

variability examined in 10 patients with an intra-class correlation coefficient of 0.99 for wall area and 0.88 for plaque components.

Conclusion

LDL lowering therapy over 2 years decreases plaque lipids without a significant change in plaque area or other components as demonstrated by MRI.

Published: 2 February 2011

doi:10.1186/1532-429X-13-S1-P382

Cite this article as: West et al.: Changes in atherosclerotic plaque composition assessed by MRI in the superficial femoral artery with two years of lipid lowering therapy. *Journal of Cardiovascular Magnetic Resonance* 2011 **13**(Suppl 1):P382.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

 BioMed Central