

ORAL PRESENTATION

Open Access

# Combined CMR and catheterization data in determining right ventricular-arterial coupling in children and adolescents with pulmonary arterial hypertension

Uyen Truong<sup>1\*</sup>, Sonali Patel<sup>1</sup>, Brian Fonseca<sup>1</sup>, Jamie Dunning<sup>2</sup>, Dunbar Ivy<sup>1</sup>, Robin Shandas<sup>1,2</sup>, Kendall Hunter<sup>2</sup>

From 17th Annual SCMR Scientific Sessions  
New Orleans, LA, USA. 16-19 January 2014

## Background

Pulmonary arterial hypertension (PAH) remains a disease with high morbidity/mortality in pediatrics. Understanding ventricular-arterial coupling, a measure of how well matched the ventricular and vascular function is, may elucidate the pathway leading to right heart failure.


## Methods

This retrospective study included subjects with PAH who had a cardiac magnetic resonance (CMR) study within 14 days of cardiac catheterization between January 2009-August 2013. The effective elastance ( $E_a$ , index of arterial load) and right ventricular maximal end-systolic elastance ( $E_{max}$ , index of contractility) were determined by a combination of CMR and hemodynamic data.  $E_a$  is defined as (mean pulmonary arterial pressure minus pulmonary capillary wedge pressure)/stroke volume.  $E_{max}$  is defined as mean pulmonary arterial pressure/end systolic volume.  $E_a/E_{max}$  ratio was derived. Additionally, a measure of non-invasive ventricular arterial coupling (assuming PWCP is insignificant, making  $E_a/E_{max}$  = end systolic volume/stroke volume) was derived from only CMR. Pulmonary vascular resistance indexed (PVRi) and pulmonary vascular reactivity, as defined by Barst criteria (decrease in mean pulmonary artery pressure of > 20%, unchanged/increased cardiac index, and decreased/unchanged pulmonary to systemic vascular resistance ratio), were also determined. Pearson correlation coefficients were calculated between PVRi and  $E_a$ ,  $E_{max}$ , and  $E_a/E_{max}$ . Receiving operating characteristic

(ROC) curve analysis determined the diagnostic value of  $E_a/E_{max}$  in predicting vascular reactivity.


## Results

Sixteen subjects were identified for inclusion with equal gender distributions. Age ranged from 3 months to 23 years (mean 11.3±7.4 years).  $E_a$  and  $E_a/E_{max}$  increased with increasing severity defined by PVRi, with  $p < 0.001$  for both.  $E_a/E_{max}$  (range 0.43-2.82) was highly correlated with PVRi ( $r = 0.92$ , 95% CI 0.79-0.97,  $p < 0.0001$ ). Non-invasively derived ventricular arterial coupling was found



**Figure 1** Regression of ventricular arterial coupling ratio ( $E_a/E_{max}$ ) and pulmonary vascular resistance indexed by reactivity. The shaded areas represent the 95% confidence interval for each regression line. The lines depict different trajectories based on reactivity, which approached, but not reach statistical significance ( $p > 0.05$ ).

<sup>1</sup>Pediatric Cardiology, Children's Hospital Colorado, Aurora, Colorado, USA  
Full list of author information is available at the end of the article



to be significantly correlated with PVRi ( $r = 0.85$ , 95% CI 0.62-0.95,  $p < 0.0001$ ), but with a lower correlation coefficient than with Ea/Emax derived from combined hemodynamic and CMR data. Regression of Ea/Emax and PVRi demonstrated differing lines when separated by reactivity, however, the lines were not significantly different (Figure 1). ROC curve analysis (Figure 2) revealed high accuracy of the Ea/Emax ratio in determining vascular reactivity. Ea/Emax of 0.85 had a sensitivity of 100% and a specificity of 80%. The area under the curve is 0.89 ( $p = 0.008$ ), suggesting good discrimination between those who were and were not reactive.

## Conclusions

Measurement of ventricular arterial coupling, Ea/Emax, in pediatrics is feasible. Pulmonary vascular non-reactivity may be due to ventricular-arterial decoupling in which ventricular contractility fails to parallel increasing afterload in severe PAH. Use of Ea/Emax may have significant prognostic implication.

## Funding

This work was supported by UL1 TR000154 from NCATS/NIH, 5R01HL114753 from NHLBI/NIH, as well as K25-094749 and K24-081506.

## Authors' details

<sup>1</sup>Pediatric Cardiology, Children's Hospital Colorado, Aurora, Colorado, USA.

<sup>2</sup>Bioengineering, University of Colorado Denver Medical Campus, Aurora, Colorado, USA.

Published: 16 January 2014

doi:10.1186/1532-429X-16-S1-O43

**Cite this article as:** Truong et al.: Combined CMR and catheterization data in determining right ventricular-arterial coupling in children and adolescents with pulmonary arterial hypertension. *Journal of Cardiovascular Magnetic Resonance* 2014 **16**(Suppl 1):O43.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

