

WORKSHOP PRESENTATION

Open Access

Efficient calculation of g-factors for CG-SENSE in high dimensions: noise amplification in random undersampling

Mehmet Akcakaya^{1*}, Tamer A Basha¹, Warren J Manning^{1,2}, Reza Nezafat¹

From 17th Annual SCMR Scientific Sessions
New Orleans, LA, USA. 16-19 January 2014

Background

SENSE [1,2] is one of the most used parallel imaging techniques. In [1], uniform undersampling was employed to efficiently reconstruct an unalised image, whereas in [2], a conjugate gradient-based method (CG-SENSE) was used for reconstruction with arbitrary trajectories. SENSE framework allows the calculation of g-factors, characterizing the noise amplification for a given k-space trajectory and coil configuration [1]. However, calculation of g-factors for arbitrary trajectories in high dimensions is time-consuming [3]. Furthermore, noise characteristics of random undersampling, used in compressed sensing, is not well-understood. In this work, we use a Monte-Carlo (MC) method for fast calculation of g-factors for CG-SENSE similar to [4,5] and apply it to random Cartesian undersampling trajectories. **Theory:** SENSE involves a pre-whitening step [1,2], thus without loss of generality, we assume white noise. SENSE reconstruction solves $\min_{\mathbf{m}} \|\mathbf{y} - \mathbf{E}\mathbf{m}\|_2$, where \mathbf{E} is the system matrix, and \mathbf{y} are the undersampled measurements. The g-factor for the k^{th} voxel is given by $g_k = \sqrt{([\mathbf{E}^*\mathbf{E}]^{-1})_{k,k} [\mathbf{E}^*\mathbf{E}]_{k,k}}$. Inverting $\mathbf{E}^*\mathbf{E}$ is not feasible in high dimensions. Instead we note the g_k corresponds to the k^{th} diagonal of the reconstruction noise covariance matrix (for normalized coil sensitivities), where $\mathbf{n}_{\text{recon}} = (\mathbf{E}^*\mathbf{E})^{-1}\mathbf{E}^*\mathbf{n}_{\text{meas}}$, and \mathbf{n}_{meas} is measurement noise with identity covariance matrix. We calculate the sample correlation matrix using a MC approach (since sample mean goes to 0), as $1/(p-1)\sum_p \mathbf{n}_{\text{recon}}^p (\mathbf{n}_{\text{recon}}^p)^*$ for p instances of $\mathbf{n}_{\text{recon}}$. Note we only calculate and store the diagonal elements of this matrix, significantly increasing efficiency.

Methods

The MC method was first verified in a numerical simulation, where the g-factor was explicitly calculated for a 2D coil configuration, to determine how many MC simulations suffice. Whole-heart imaging was performed with an isotropic resolution of 1.3 mm using a 32-channel coil array. Two 4-fold accelerated acquisitions were performed, one with uniform undersampling (2×2 in the k_y - k_z plane) and one with random undersampling. Coil sensitivity maps were exported. Images were reconstructed using SENSE (for uniform) and CG-SENSE (for both). g-factors were also calculated with the proposed approach.

Results

Figure 1 shows the results of numerical simulations, indicating the method converges in ~ 50 MC simulations. Figure 2 shows the reconstructions associated with the two undersampling patterns and reconstructions, and the corresponding g-factors respectively. The results exhibit the semi-convergence property for random undersampling but not for uniform. Furthermore, the g-factor for random undersampling is smaller at its convergent point than for uniform.

Conclusions



g-factors for random undersampling is better than those for uniform at high k-space dimensions and high acceleration rates.

Funding

NIH:K99HL111410-01; R01EB008743-01A2.

¹Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, USA

Full list of author information is available at the end of the article



Authors' details

¹Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, USA. ²Radiology, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, USA.

Published: 16 January 2014

References

1. Pruessmann: *MRM* 1999.
2. Pruessmann: *MRM* 2001.
3. Liu: *ISMRM* 2008.
4. Thunberg: *MRI* 2007.
5. Robson: *MRM* 2008.

doi:10.1186/1532-429X-16-S1-W28

Cite this article as: Akcakaya et al.: Efficient calculation of g-factors for CG-SENSE in high dimensions: noise amplification in random undersampling. *Journal of Cardiovascular Magnetic Resonance* 2014 16(Suppl 1):W28.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

